

Crossover from random-exchange to random-field critical behavior in Ising models.

Pasquale Calabrese,¹ Andrea Pelissetto,² Ettore Vicari³

¹ *Scuola Normale Superiore and INFN, Piazza dei Cavalieri 7, I-56126 Pisa, Italy.*

² *Dipartimento di Fisica dell'Università di Roma I and INFN, I-00185 Roma, Italy.*

³ *Dipartimento di Fisica dell'Università and INFN, Via Buonarroti 2, I-56127 Pisa, Italy.*

e-mail: calabres@df.unipi.it, Andrea.Pelissetto@roma1.infn.it, vicari@df.unipi.it.

(February 2, 2008)

Abstract

We compute the crossover exponent ϕ describing the crossover from the random-exchange to the random-field critical behavior in Ising systems. For this purpose, we consider the field-theoretical approach based on the replica method, and perform a six-loop calculation in the framework of a fixed-dimension expansion. The crossover from random-exchange to random-field critical behavior has been observed in dilute anisotropic antiferromagnets, such as $\text{Fe}_x\text{Zn}_{1-x}\text{F}_2$ and $\text{Mn}_x\text{Zn}_{1-x}\text{F}_2$, when applying an external magnetic field. Our result $\phi = 1.42(2)$ for the crossover exponent is in good agreement with the available experimental estimates.

PACS Numbers: 75.10.Nr, 75.10.Hk, 64.60.Ak

The most studied experimental realizations of random-field Ising systems are dilute anisotropic antiferromagnets in a uniform magnetic field applied along the spin ordering axis [1–3]. A simple lattice model is provided by the Hamiltonian

$$\mathcal{H} = J \sum_{\langle ij \rangle} \rho_i \rho_j s_i s_j - H \sum_i \rho_i s_i, \quad (1)$$

where $J > 0$, the first sum extends over all nearest-neighbor sites, $s_i = \pm 1$ are the spin variables, and ρ_i are uncorrelated quenched random variables, which are equal to one with probability p (the spin concentration) and zero with probability $1 - p$ (the impurity concentration). In the absence of an external field, i.e. $H = 0$, the critical behavior of dilute Ising systems (above the percolation point of the spins) belongs to the universality class of the random-exchange Ising model (REIM), which differs from the standard Ising one. See, e.g., Refs. [4,5] for recent reviews. The applied uniform field H gives rise to a different critical behavior corresponding to the universality class of the random-field Ising model (RFIM). See e.g. Refs. [6–8] for recent reviews on RFIM.

For $t \equiv (T - T_N)/T_N \rightarrow 0$, T_N being the Néel temperature in zero field, and $H \rightarrow 0$ the singular part of the free energy can be written as [1]

$$\mathcal{F} = |u_t|^{2-\alpha} f(H^2 |u_t|^{-\phi}), \quad (2)$$

where $u_t \approx t + a_1 H^2 + a_2 t^2$ is the scaling field associated with the temperature, α is the specific-heat exponent of the REIM, $f(x)$ is a scaling function, and ϕ is the corresponding crossover exponent. As a consequence of the crossover scaling (2), the critical temperature in the presence of the external field H is given by

$$T_c(H) \approx T_N + c H^{2/\phi} - a H^2, \quad (3)$$

for sufficiently small H . Experimental measurements on dilute antiferromagnets yielded rather precise estimates: $\phi = 1.42(3)$ obtained for $\text{Fe}_x\text{Zn}_{1-x}\text{F}_2$ [9], $\phi = 1.43(3)$ for $\text{Mn}_x\text{Zn}_{1-x}\text{F}_2$ [10], and $\phi = 1.41(5)$ for $\text{Fe}_x\text{Mg}_{1-x}\text{Cl}_2$ [11].

On the theoretical side, a computation of ϕ was presented in Ref. [12] using the field-theoretical ϵ expansion. We recall that in the REIM the ϵ expansion is actually an expansion in powers of $\sqrt{\epsilon}$. The exponent ϕ was computed to $O(\epsilon)$ [12], obtaining

$$\phi/\gamma = 1 + 0.16823\epsilon^{1/2} - 0.22666\epsilon + O(\epsilon^{3/2}), \quad (4)$$

where γ is the REIM susceptibility exponent. Apart from showing that $\phi \neq \gamma$, where $\gamma \approx 1.34$ (see, e.g., Refs. [13–15]), this series does not yield a reliable estimate of ϕ for three-dimensional systems. This is generically true for the $\sqrt{\epsilon}$ expansion of any critical exponent of the REIM [4,5].

In this paper we determine the crossover exponent ϕ using an alternative field-theoretical method based on a fixed-dimension expansion in powers of appropriate zero-momentum quartic couplings, which we compute to six loops. As we shall see, the analysis of the series provides a quite precise estimate of ϕ ,

$$\phi = 1.42(2), \quad (5)$$

in good agreement with experiments. This result was already anticipated in Ref. [16].

The field-theoretical approach is based on an effective Landau-Ginzburg-Wilson φ^4 Hamiltonian that can be obtained by using the replica method [17], i.e.

$$\mathcal{H}_{\varphi^4} = \int d^3x \left\{ \frac{1}{2} \sum_{i=1}^N [(\partial_\mu \varphi_i)^2 + r \varphi_i^2] + \frac{1}{4!} \sum_{i,j=1}^N (u_0 + v_0 \delta_{ij}) \varphi_i^2 \varphi_j^2 \right\}, \quad (6)$$

where φ_i is an N -component field. The critical behavior of the REIM is expected to be described by the stable fixed point of the Hamiltonian $\mathcal{H}_{\varphi^4}$ in the limit $N \rightarrow 0$ for $u_0 < 0$ [18]. The most precise field-theoretical results for the critical exponents have been obtained by analyzing the fixed-dimension expansion in powers of the zero-momentum quartic couplings u, v related to u_0, v_0 (i.e. $u = Z_u u_0/m$ and $v = Z_v v_0/m$ where $Z_{u,v} = 1 + O(u, v)$), which have been computed to six loops [15,19]. We refer the reader to Ref. [15] for notations and definitions.

In the presence of a spatially uncorrelated random field $h(x)$ with zero average and variance h^2 , one may still apply the replica method. Averaging the replicated random field term

$$\int d^3x h(x) \sum_i \varphi_i(x) \quad (7)$$

over the Gaussian distribution of the field, one obtains a new term proportional to

$$h^2 \int d^3x \sum_{i,j} \varphi_i(x) \varphi_j(x), \quad (8)$$

that must be added to the Hamiltonian $\mathcal{H}_{\varphi^4}$ [12]. This term causes the crossover from the REIM to the RFIM. Thus, the REIM-to-RFIM crossover exponent is determined by the $N \rightarrow 0$ limit of the renormalization-group (RG) dimension y_T of the quadratic operator [20]

$$T_{ij} = \varphi_i \varphi_j, \quad i \neq j. \quad (9)$$

In order to evaluate y_T and the corresponding crossover exponent $\phi = y_T \nu$, we define a related RG function Z_T from the one-particle irreducible two-point function $\Gamma_T^{(2)}$ with an insertion of the operator T_{ij} , i.e.

$$\Gamma_T^{(2)}(0)_{ij,kl} = Z_T^{-1} A_{ijkl}, \quad (10)$$

where $A_{ijkl} = \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}$ so that $Z_T(0) = 1$. Then, we compute the RG function

$$\eta_T(u, v) = \left. \frac{\partial \ln Z_T}{\partial \ln m} \right|_{u_0, v_0} = \beta_u \frac{\partial \ln Z_T}{\partial u} + \beta_v \frac{\partial \ln Z_T}{\partial v}, \quad (11)$$

where β_u and β_v are the β -functions. The exponent η_T is given by the value of $\eta_T(u, v)$ at $u = u^*$, $v = v^*$, where (u^*, v^*) is the REIM stable fixed point. Finally, the REIM-to-RFIM crossover exponent is obtained by using the RG scaling relation

$$\phi = (2 + \eta_T - \eta) \nu = \gamma + \eta_T \nu, \quad (12)$$

where γ , ν , and η are REIM critical exponents.

We computed the function $\Gamma^{(T,2)}(0)$ to six loops. The calculation is rather cumbersome, since it requires the evaluation of 563 Feynman diagrams. We handled it with a symbolic manipulation program, which generates the diagrams and computes the symmetry and group factors of each of them. We used the numerical results compiled in Ref. [21] for the integrals associated with each diagram. The resulting six-loop series of $\eta_T(u, v)$ is

$$\begin{aligned} \eta_T(\bar{u}, \bar{v}) = & -\frac{1}{4}\bar{u} + \frac{1}{16}\bar{u}^2 + \frac{1}{18}\bar{u}\bar{v} - 0.0357673\bar{u}^3 - 0.0483240\bar{u}^2\bar{v} - 0.00421548\bar{u}\bar{v}^2 \\ & + 0.0343748\bar{u}^4 + 0.0762616\bar{u}^3\bar{v} + 0.0416943\bar{u}^2\bar{v}^2 + 0.00896116\bar{u}\bar{v}^3 \\ & - 0.0408958\bar{u}^5 - 0.121377\bar{u}^4\bar{v} - 0.104778\bar{u}^3\bar{v}^2 - 0.0373712\bar{u}^2\bar{v}^3 - 0.00527015\bar{u}\bar{v}^4 \\ & + 0.0597048\bar{u}^6 + 0.227662\bar{u}^5\bar{v} + 0.287108\bar{u}^4\bar{v}^2 + 0.17231\bar{u}^3\bar{v}^3 \\ & + 0.0552124\bar{u}^2\bar{v}^4 + 0.00759654\bar{u}\bar{v}^5 + \dots, \end{aligned} \quad (13)$$

where $\bar{u}$ and $\bar{v}$ are the rescaled couplings $\bar{u} = u/(6\pi)$ and $\bar{v} = 3v/(16\pi)$, and the dots indicate higher-order terms. In our analysis we also considered the series $\phi(\bar{u}, \bar{v})$ corresponding to ϕ , which can be obtained by using Eq. (12) and the series of the RG functions $\gamma(\bar{u}, \bar{v})$ and $\nu(\bar{u}, \bar{v})$ corresponding to the critical exponents γ and ν , cf. Ref. [15].

In order to obtain an estimate of η_T and ϕ , the corresponding six-loop series must be resummed and then evaluated at the fixed-point values u^* , v^* . Although the perturbative expansion is not Borel summable [22], various resummation schemes have been proposed and employed, obtaining rather reliable results for the critical exponents, see, e.g., Refs. [4,5]. We analyzed $\eta_T(\bar{u}, \bar{v})$ and $\phi(\bar{u}, \bar{v})$ by using the resummation methods outlined in Ref. [15] and the estimates $u^* = -18.6(3)$ and $v^* = 43.3(2)$ obtained by Monte Carlo simulations in Ref. [13], which turn out to be more precise than those obtained from the zeroes of the β -functions computed to six loops.

We obtained $\eta_T = 0.095(30)$ from the analysis of $\eta_T(\bar{u}, \bar{v})$, $\phi = 1.43(3)$ from $\phi(\bar{u}, \bar{v})$, and $\phi = 1.42(2)$ from $1/\phi(\bar{u}, \bar{v})$. The errors are related to the spread of the results obtained by using different resummation methods and different resummation parameters within each method, see Ref. [15] for details. Using the available estimates of the critical exponents (e.g., $\gamma = 1.342(6)$, $\nu = 0.683(3)$ and $\eta = 0.035(2)$ from Monte Carlo simulations [13,14], $\gamma = 1.330(17)$, $\nu = 0.678(10)$ and $\eta = 0.030(3)$ from the analysis of the six-loop field-theoretical expansions [15]), the above-reported result for η_T and the RG scaling relation (12) give $\phi = 1.41(2)$. From these results we arrive at the final estimate reported in Eq. (5), which is in good agreement with the available experimental results obtained for various uniaxial anisotropic antiferromagnets.

REFERENCES

- [1] S. Fishman and A. Aharony, J. Phys. C **12**, L729 (1979).
- [2] P. Z. Wong, S. von Molnar, and P. Dimon, J. Appl. Phys. **53**, 7954 (1982).
- [3] J. L. Cardy, Phys. Rev. B **29**, 505 (1984).
- [4] A. Pelissetto and E. Vicari, Physics Reports **368**, 549 (2002), arXiv:cond-mat/0012164.
- [5] R. Folk, Yu. Holovatch, and T. Yavors’kii, Uspekhi Fizicheskikh Nauk **173**, 175 (2003) [Physics Uspekhi **46**, 175 (2003)].
- [6] D. P. Belanger, in *Spin Glasses and Random Fields*, edited by A. P. Young (World Scientific, Singapore, 1998), p. 251.
- [7] T. Nattermann, in *Spin Glasses and Random Fields*, edited by A. P. Young (World Scientific, Singapore, 1998), p. 277.
- [8] D. P. Belanger, Brazilian J. Phys. **30**, 682 (2000).
- [9] I. B. Ferreira, A. R. King, and V. Jaccarino, Phys. Rev. B **43**, 10797 (1991); J. Appl. Phys. **69**, 5246 (1991).
- [10] C. A. Ramos, A. R. King, and V. Jaccarino, Phys. Rev. B **37**, 5483 (1988).
- [11] U. A. Leitão and W. Kleemann, Phys. Rev. B **35**, 8696 (1987).
- [12] A. Aharony, Europhys. Lett. **1**, 617 (1986).
- [13] P. Calabrese, V. Martín-Mayor, A. Pelissetto, and E. Vicari, arXiv:cond-mat/0306272.
- [14] H. G. Ballesteros, L. A. Fernández, V. Martín-Mayor, A. Muñoz Sudupe, G. Parisi, and J. J. Ruiz-Lorenzo, Phys. Rev. B **58**, 2740 (1998).
- [15] A. Pelissetto and E. Vicari, Phys. Rev. B **62**, 6393 (2000).
- [16] P. Calabrese, A. Pelissetto, P. Rossi, and E. Vicari, Talk at the International Conference on Theoretical Physics, TH-2002, UNESCO, Paris, 2002, hep-th/0212161.
- [17] V. J. Emery, Phys. Rev. B **11**, 239 (1975); S. W. Edwards and P. W. Anderson, J. Phys. F **5**, 965 (1975); G. Grinstein and A. Luther, Phys. Rev. B **13**, 1329 (1976).
- [18] In a recent work [P. Calabrese, P. Parruccini, A. Pelissetto, and E. Vicari, in preparation] we showed that the attraction domain of the REIM fixed-point contains at least the region $-1 \lesssim u_0/v_0 < 0$.
- [19] J. M. Carmona, A. Pelissetto, and E. Vicari, Phys. Rev. B **61**, 15136 (2000).
- [20] In the cubic-symmetric theory (6) the RG dimension of the quadratic operators $U_i = \varphi_i^2 - \frac{1}{N} \sum_a \varphi_a^2$ is different from the one of $T_{ij} = \varphi_i \varphi_j$ with $i \neq j$, unlike $O(N)$ -symmetric theories where they belong to the same spin-2 irreducible representation. In the limit $N \rightarrow 0$ the RG dimensions of U_i and the invariant operator $\frac{1}{N} \sum_a \varphi_a^2$ turn out to be equal. This can be inferred from their perturbative expansions.
- [21] B. G. Nickel, D. I. Meiron, and G. A. Baker, Jr., “Compilation of 2-pt and 4-pt graphs for continuum spin models,” Guelph University Report, 1977, unpublished.
- [22] A. J. Bray, T. McCarthy, M. A. Moore, J. D. Reger, A. P. Young, Phys. Rev. B **36**, 2212 (1987); A. J. McKane, Phys. Rev. B **49**, 12003 (1994); G. Álvarez, V. Martín-Mayor, J. J. Ruiz-Lorenzo, J. Phys. A **33**, 841 (2000).